

Final Exam MAT 295 Fall 2002

To receive full or partial credit the correct work leading to the correct answer must be written down.

Your Name	Instructor's Name

Your Signature:

Problem	Score
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
Total	

2

(1) Compute the limits.

(a) (4pts) $\lim_{x \rightarrow 0} \frac{1 - \cos(x)}{\sin(x)}$

(b) (4pts) Given $3 \leq f(x) \leq x^2 + 6x + 12$ find $\lim_{x \rightarrow -3} f(x)$

(2) (6pts) Determine the constant c so that $f(x)$ is continuous at $x = 1$

where $f(x) = \begin{cases} \ln(x) & 0 < x \leq 1 \\ e^{x-1} + c & x > 1 \end{cases}$

(3) Differentiate **do not simplify**

(a) (4pts) Find $\frac{d}{ds} \frac{3s^{-1} + 6}{2s^{1/2} + 3}$.

(b) (4pts) $r(\theta) = \sin(\cos(\theta^2))$. Find $\frac{d}{d\theta} r(\theta)$.

(c) (4pts) Use the properties of logarithms first and then differentiate to find $\frac{d}{dx} \ln \left(\frac{\sin^4(x)}{x^2(x+5)^{10}} \right)$

(d) (4pts) Find $\frac{d}{dx} \int_0^{2x} \frac{t^2 - 4}{1 + \cos^2(t)} dt$

(e) (4pts) Find $\frac{d}{dx} (\sin(x))^x$

4

(4) (5pts) Find the slope of the tangent line to the curve $x^2y^2 + x^3 + y^3 = 3$ at the point $(1, 1)$

(5) (5pts) The circumference of a sphere is measured to be 86 cm with a maximum possible error of 0.5cm. Estimate the relative error in calculating the volume of the sphere ($V = \frac{4}{3}\pi r^3$) with this measurement.

- (6) (5pts) The top half of an hour glass is an inverted cone with height 4 cm and **diameter** 2 cm. The volume of a cone is $V = \frac{1}{3}\pi r^2 h$. Sand is running out of this cone at the rate of $.01 \text{ cm}^3/\text{sec}$. How fast is the depth of sand in this cone decreasing when the depth is 3 cm?
- (7) (6pts) A rectangle is inscribed with its base on the x -axis and its upper corners on the top half of the circle $y = \sqrt{4 - x^2}$. What is the largest possible area for such a rectangle? Be sure to show that your answer is a maximum.

(8) You are given the function $f(x) = \frac{(x-1)^3}{x^3}$ along with its first derivative $f'(x) = \frac{3(x-1)^2}{x^4}$ and second derivative $f''(x) = -6 \frac{(x-1)(x-2)}{x^5}$.

- (a) (2pts) Find all vertical asymptotes. You must take the appropriate limits to receive credit.
- (b) (2pts) Find all horizontal asymptotes. You must take the appropriate limits to receive credit.
- (c) (2pts) Find all critical points. Show your work.
- (d) (3pts) List all intervals on which f is increasing. Show your work.
- (e) (3pts) List all intervals on which f is concave down. Show your work.
- (f) (2pts) Find all inflection points. Show your work.

- (9) (6pts) Calculate the Riemann sum R_n for $f(x) = 3x^2 + 1$ on the interval $[0, 3]$ by dividing the interval into n equal width subintervals and using right hand endpoints as sample points.
- (10) (5pts) On the interval $[0, \pi]$ the function $f(x) = \cos(x)$ is one-to-one, so it has an inverse function $f^{-1}(x)$ whose domain is $[-1, 1]$. Find $(f^{-1})'(0)$, i.e. the derivative of the inverse function at 0.

(11) Compute the definite and indefinite integrals.

(a) (4pts) Given that $\int_1^{10} f(x)dx = 8$, $\int_1^4 f(x)dx = -2$, and $\int_8^{10} f(x)dx = 5$

find $\int_4^8 f(x)dx$.

(b) (4pts) $\int_1^4 \frac{2x^2 + 7}{\sqrt{x}} dx$.

(c) (4pts) $\int_0^{\pi/3} \cos(3\theta) d\theta$.

(d) (4pts) $\int \frac{x}{\sqrt{4x^2 - 8}} dx$.

(e) (4pts) $\int e^{4x} + \frac{3}{x} dx$.